RTC Dental –

<u>Instructions for Laser Perio Therapy:</u>

• Please read through these heath review questions and send RTC Dental a list of conditions you suffer from:

```
(Email <u>info@RTCdental.com</u>)
(Dr. Ali's Cell Phone #: 571-423-7945)
```

- O What is the **phone number for your preferred pharmacy**, so our dentist can call in your prescriptions for you?
 - What is your full name and date of birth?
- o Allergies:
 - Are you allergic to penicillin or amoxicillin antibiotics?
 - Are you allergic to pain killers such as hydrocodone or Vicodin?
- o Illnesses:
 - Do you have heart problems or any major illness that your dentist should be aware of?
 - Do you have any joint replacement surgeries in your past?
 - Hip replacement surgery?
 - Knee replacement surgery?
 - Elbow replacement surgery?
 - Do you have a pace maker?
 - Do you have history of stroke, heart attack, angioplasty or heart surgery?
 - Do you have any bleeding disorders?
 - Do you have Diabetes?
 - If so, then what are your typical blood sugar levels when you are tested?
 - Do you have hypertension?
 - If so, then what are your typical blood pressure numbers when you are tested?
 - Do you ever have difficulty climbing stairs, or become winded when climbing stairs?
 - What kind of physical activities do you typically do?

Special Pre-Op Medicines:

- Rinse with Chlorhexidine in advance of your laser session:
 - Rinse and spit with chlorhexidine mouth rince, 2-3 times daily to kill bacteria
- Take Antibiotic Pills in advance of your Laser Perio session:
 - Penicillin or Clindamycin antibiotics
 - Start taking antibiotic pills ~24 hours prior to your 1st laser session.
 - Continue taking your antibiotic pills for 10 full days.
 - We recommend you complete your 2nd laser session prior to the end of your antibiotic prescription.
 - Only stop taking the antibiotic pills if you experience allergic reaction or GI disturbance.
 - Allergic reaction to antibiotics typically involves rash on skin; and in severe cases can lead to difficulty breathing, if your throat swells.
 - O GI disturbance are typically diarrhea; but in more severe cases may involve bloody stools.
 - Adverse reactions to antibiotics pills should be reported to your dentist immediately.
 - Call Dr. Ali Miamee's cell phone number 571-423-7945
 - If you are experiencing difficulty breathing or swallowing spit, then you should immediately visit an emergency room or call 911.

<u>Instructions For The Day Of The Procedure:</u>

- On the day of your procedure please do the following:
 - Eat a large rich breakfast to ensure there is plenty of sugar in your blood during your appointment.
 - Eggs, meats, sausage, butter, syrups are all preferred over fruits or vegetables.
 - High calorie foods are preferred.
 - o Brush your teeth well before leaving for your appointment.
 - You will be asked not to brush your teeth for 24 hours after your appointment.
 - Rinse with chlorhexidine prior to leaving for your laser session.
 - Take all normal medications on schedule, as prescribed from your physicians.
 - Hypertension meds
 - Diabetes meds
 - Blood thinners

- Feel free to call (or text) Dr. Ali Miamee on his cell phone with any questions:
 - o Cell phone # 571-423-7945

<u>Instructions For After Your Procedure:</u>

- Post-Op Instructions for after each laser perio surgery:
 - Do not brush your teeth for 24 hours after the procedure.
 - Gentle brushing of your teeth can resume after 24 hours.
 - Do not floss your teeth for 1-2 weeks after the procedure.
 - Gentle flossing can resume after 1 week.
 - Normal flossing can resume after 2 weeks.
 - Do not brush your gums for 1-2 weeks after the procedure.
 - Gentle brushing of the gums can resume after 1 week.
 - Normal brushing of the gums can resume after 2 weeks.
 - Continue using chlorhexidine mouth rinse for 1-2 weeks after your procedure.
 - Stop using chlorhexidine and call your dentist if you see discoloration on your teeth due to chlorhexidine use.
 - Schedule a follow up appointment with your dentist for 7-14 days postop, to evaluate your healing.